

## **Atopic Dermatitis** Patient Enrollment and Prescription Form

| PATIENT INFORMATION  | PRESCRIBER INFORMATION                         |  |  |
|--|--|--|--|
| Patient Name   | Prescriber's Name                              |  |  |
| DOB/   |  |  |  |
| Weight Height Phone  |  |  |  |
| Address  |  |  |  |
| City, State, Zip  Cell Phone E-Mail  | City, State, Zip Fax Fax                       |  |  |
| PLEASE PROVIDE COPY OF PATIENT'S PRESCRIPTION INSURANCE CARD WITH ENROLLMENT FORM  |  | Office Contact Person  |  |
| Insurance Co. Name   |  |  |  |
| Insurance Co. Phone Group#   | <del></del>                                    |  |  |
| insurance co. Prione Group#  | With my signature on this form, I also authori |  |  |
| Policy Holder Name   | which includes serving as my prior authorizati | on designated agent in dealing with  |  |
| Policy Holder Employer   |  | A Company of the Comp |  |
| Relationship to Patient  |  |  |  |
| ID# RxBIN PCN  | Ship to: □Patient □Prescriber's Off            |  |  |
|  | CHART NOTES FOR INSURANCE PUR                  | POSES  |  |
| Medical Information (DO NOT COMPLETE IF CHART NOTES ARE  |  |  |  |
| Diagnosis ICE  | 4 300 5  | (A)  |  |
| Date of Diagnosis Body Surface Affected  |  | M IT   |  |
| <ul> <li>Patient has moderate to severe atopic dermatitis (AD) that is<br/>controlled on prior or current topical therapy</li> </ul> | sinadequately                                  |  |  |
|  |  |  |  |
| Drug Allergies Latex A   | Allergy? DYES DNO                              | 1 / Suffer what  |  |
| Failed Prior therapies?  | 1// //   | 1 11 17 12 11  |  |
| ☐ Topical Corticosteroids:   | 2/1/2/1  | 2 5/1/1/1/   |  |
| ·  | 054/0 \ /\ /\                                  | <b>歌                                    </b>   |  |
|  | to   | (3)  |  |
| ☐ Topical therapy not appropriate  | 1.77/1.1                                       |  |  |
| Reason:  | V 1817   | ( )( )   |  |
| ☐ Systemic Corticosteriods, Immunosupressants and/or P   | 1 0 /  | 14 111   |  |
|  | 7-11 V   | 17 1359  |  |
|  |  | 23 99  |  |
| ☐ Systemic corticosteroids not appropriate   |  | ☐ Hands ☐ Feet☐ Nails ☐ Face   |  |
| ☐ Immunosuppressants not appropriate   | Other:   |  |  |
| ☐ Phototherapy not appropriate   |  |  |  |
| Reason:  |  |  |  |
| Does the patient have history of conjunctivitis or keratitis?   YES  Additional Comments   | □NO  |  |  |
| PRESCRIPTION INFORMATION   | (please indicate days suppl                    | y if different than suggested days supply)   |  |
| □Adbry® (tralokinumab-ldrm) □ 150mg Pre-filled Syri  |  |  |  |
| □ ADULT DOSING: 600MG SQ at week 0 and 300MG   | every other week Quantity:                     | Refills  |  |
| □ PEDIATRIC DOSING: 300MG at week 0 and 150MG  |  | Refills  |  |
| Tag  | mg Tablet                                      |  |  |
| Directions: Take 1 tablet daily  | Qty: 30 tablets                                | Refills  |  |
| □ Dupixent® (dupilumab) □ Syringe □ Pen  |  |  |  |
| □ ADULT STARTER: 600MG SQ at week 0, 300MG S   |  | Refills  |  |
| □ ADULT MAINTENANCE: 300MG SQ once every 2 w   |  |  |  |
| □ PED (up to 5yr): 200MG SQ every 4 weeks  | Quantity: 2                                    | <del></del>  |  |
|  |  |  |  |
| □ PED (up to 5yr): 300MG SQ every 4 weeks  | Quantity: 2                                    |  |  |
| □ PED (6-17 yrs) up to 30kg: <u>600MG_SQ_week 0, 300</u>   |  |  |  |
| □ PED (6-17 yrs) up to 60kg: <u>400MG_SQ week 0, 200</u>   |  |  |  |
| ☐ PED (6-17 yrs) above 60kg: <u>600MG_SQ</u> week 0, 30  | 00mg Q2 weeks Quantity: 2                      | Refills  |  |
| □ EBGLYSS™ (lebrikizumab-lbkz) □ 250mg/2mL Pre-F   | illed Syringe 🔲 250mg/2mL Pen                  |  |  |
| ☐ INITIAL DOSE: <u>500MG SQ at week 0 and week 2</u>   | Quantity:                                      | Refills  |  |
| ☐ INDUCTION DOSE: _250 MG SQ every 2 weeks   | Quantity:                                      | Refills  |  |
| ■ MAINTENANCE DOSE: 250MG SQ every 4 weeks   | Quantity:                                      | Refills  |  |
| ☐ OPZELURA™ 1.5% Cream 60 GM Tube  |  |  |  |
| Directions: Apply thin layer twice daily to affected are   | as (up to 20% BSA) MAX 60 gm/wk. Qt            | y:tube(s) Refills  |  |
| □ Rinvoq® (upadacitinib) □ 15mg ER Tablet □ 30m  | ng ER Tablet                                   |  |  |
| Directions:Take 1 tablet daily   | Qty: 30 tablets                                | Refills  |  |
| □VTAMA® (tapinarof) cream 60 gram tube   |  |  |  |
| Directions: Apply a thin layer of VTAMA cream to affected  | d areas once daily Quantity                    | Tube(s) Refills  |  |