

Crohn's Disease / Ulcerative Colitis Patient Enrollment & Prescription Form

rel: 844-443-6879	PRESCRIBER INFORMATION
Patient Name	Prescriber's Name
DOB/ SSN Gender	Practice Name
Weight Height Phone	
Address	
City, State, Zip	
Cell PhoneE-Mail PLEASE PROVIDE COPY OF PATIENT'S PRESCRIPTION INSURANCE CARD WITHENROLLMENT FORM	Phone Fax Office Contact Person
Insurance Co. Name	
Insurance Co. Name Group#	
	With my signature on this form, Lake authorize use of Centry Health's Services
Policy Holder Name Policy Holder Employer	
Relationship to Patient	Described of the control of the cont
ID# RxBIN PCN_	
	CHART NOTES FOR INSURANCE PURPOSES
Medical Information (DO NOT COMPLETE IF CHART NOTES A	ARE PROVIDED)
Diagnosis	Date of Diagnosis
CD-10 Code: New to therapy? \Box	YES NO
	tis B negative?
	Latex Allergy? YES NO
Prior Therapy and for how long?	
Reason for Discontinuation?	
Any other relevant medical info?	
PRESCRIPTION INFORMATION (DAW requests must be handwritten) ☐ Cimzia® (certolizumab) ☐ 200mg/1ml Prefilled Syringe St	cartor Vit
Directions: Inject 400 mg SQ initially and at Weeks 2 and 4	
☐ 200mg/1ml Prefilled Syringe fo	
Directions: ☐ Inject 400 mg SQ every 4 wks OR ☐ Inject	ct 400 mg SQ every 2 wks
	write DAW):
□ Crohn's 40mg Starter Package	o weeks later (Day 15) Qty:1 KIT Refills0
□ 40mg Prefilled Pen Carton for Maintenance Dosing	o weeks later (Day 15) Qty 1 Kii Kelliis
Directions: Inject 40mg every other week	Qty: <u>QS 30 Days</u> Refills
□ ENTYVIO® (vedolizumab) 108mg/0.68mL Pen	Qty: 2 Pens Refills
Directions: <u>Inject 108mg subcutaneously every 2 weeks</u> □ OMVOH® (mirikizumab) □ INDUCTION DOSE □ 300 mg	via I.V. at Week 0, Week 4, and Week 8 Qty: 1 vial Refills zero
• • •	via I.V. at Week 0, Week 4, and Week 8 Qty: 1 vial Refills <u>zero</u> SQ at Week 12 & every 4 weeks thereafter Qty: 2 PFS Refills
(given as	two consecutive injections of 100 mg each)
· · · · <u> · </u>	ne 45mg tab once daily for 8 weeks
	ne 15mg tablet once daily Qty: 30 Refills ne 30mg tablet once daily Qty: 30 Refills
☐ Simponi® (golimumab) TYPE: ☐ Smartject® Autoinjector ☐	
☐ 15 - 39kg Dosing & Directions: 100 mg SQ initially Week 0, 5	0 mg at Week 2, then 50 mg every 4 weeks Qty: QS 30 Days Refills
□ ≥ 40kg Dosing & Directions: 200 mg SQ initially Week 0, 100	
□ Skyrizi® (risankizumab-rzaa) □ 600 mg single-dose vial - indu □ INITIATION: Infuse 600 mg as initial IV dose at Week 0, W	uction 180mg single dose cartridge 360mg single dose cartridge Qty: 60 Days
☐ MAINTENANCE: 180 mg by SQ injection at week 12, and every state of the state of	•
☐ MAINTENANCE: 360 mg by SQ injection at week 12, and ex	
	90mg single-dose prefilled syringe
 260mg Starter (up to 55kg): <u>Infuse intravenously over a p</u> 390mg Starter (greater than 55kg to 85kg): Infuse intrave 	
☐ 520mg Starter (greater than 85mg): Infuse intravenously	
☐ MAINTENANCE: <u>Inject 90mg SQ every 8 weeks after initial</u>	al intravenous dose 60 Days Supply? □YES □NO Refills
□ Tremfya® (guselkumab) □ 100mg Pen □ 100mg Prefilled S	
□ INITIATION (Crohn's Only): Inject 400mg SQ at Week 0, Week 10, and every 8	<u> </u>
☐ MAINTENANCE: Inject 200mg SQ at Week 12 and every 4 v	
- · · · · · · · · · · · · · · · · · · ·	ne 10mg tab twice daily for 8 weeks Qty: 120
· · · · · · · · · · · · · · · · · · ·	ne 22mg XR tab once daily for 8 weeks Qty: 30
	ne 5mg tablet twice daily Qty: 60 Refills ne 10mg tablet twice daily Qty: 60 Refills
· · · · · · · · · · · · · · · · · · ·	ne 11mg XR tablet once daily
□ <u>Take o</u>	ne 22mg XR tablet once daily Qty: 30 Refills
□ Zeposia® (ozanimod) □ 7 Day Starter □ 0.92mg Ca	
☐ INDUCTION DOSE: Take 0.23 mg once daily for days 1-4, the	en take 0.46 mg once daily for days 5-7 Qty: 1 Pack