Crohn's Disease / Ulcerative Colitis
Patient Enrollment & Prescription Form

Tel: 844-443-6879 **Fax: 844-329-2447** ePrescribe to our pharmacy at "GENTRY HEALTH SERVICES" in Avon Lake, Ohio.

PATIENT INFORMATION

Patient Name _____
DOB ___/___/_____ SSN _____ Gender_____
Weight _____ Height _____ Phone _____
Address _____
City, State, Zip _____
Cell Phone _____ E-Mail _____

PLEASE PROVIDE COPY OF PATIENT'S PRESCRIPTION INSURANCE CARD WITH ENROLLMENT FORM

Insurance Co. Name _____
Insurance Co. Phone _____ Group# _____
Policy Holder Name _____
Policy Holder Employer _____
Relationship to Patient _____
ID# _____ RxBIN _____ PCN_____

PRESCRIBER INFORMATION

Prescriber's Name _____
Practice Name _____
DEA _____ NPI _____
Address _____
City, State, Zip _____
Phone _____ Fax _____
Office Contact Person _____
Office Contact EMAIL _____
Prescription Date _____ Date Needed _____

With my signature on this form, I also authorize use of Gentry Health's Services which includes serving as my prior authorization designated agent in dealing with medical and prescription insurance companies, and co-pay assistance foundations.

Prescriber Signature _____

Ship to: ☐Patient ☐Prescriber's Office ☐Discount Drug Mart

PLEASE PROVIDE ALL RELEVANT CHART NOTES FOR INSURANCE PURPOSES

Medical Information (DO NOT COMPLETE IF CHART NOTES ARE PROVIDED)

Diagnosis _____ Date of Diagnosis _____
ICD-10 Code: _____ New to therapy? ☐ YES ☐ NO
TB negative? ☐ YES ☐ NO Hepatitis B negative? ☐ YES ☐ NO
Drug Allergies _____ Latex Allergy? ☐ YES ☐ NO
Prior Therapy and for how long? _____
Reason for Discontinuation? _____
Any other relevant medical info? _____

PRESCRIPTION INFORMATION *(DAW requests must be handwritten)*

☐ **Cimzia® (certolizumab)** ☐ 200mg/1ml Prefilled Syringe Starter Kit
Directions: Inject 400 mg SQ initially and at Weeks 2 and 4 Qty: 6 Refills 0
☐ 200mg/1ml Prefilled Syringe for Maintenance Dosing
Directions: ☐ Inject 400 mg SQ every 4 wks **OR** ☐ Inject 400 mg SQ every 2 wks Qty: QS 30 Days Refills _____

☐ **adalimumab** ☐ Prefilled Syringe ☐ Prefilled Pen ☐ **Please check if Starter Dose is NOT needed**
☐ **Preferred Brand Name** (if required must write DAW): _____
Directions: 160mg SQ on day 1, 80mg on day 15, then 40mg every other week starting day 29 Quantity _____ Refills _____

☐ **ENTYVIO® (vedolizumab)** 108mg/0.68mL Pen Qty: 2 Pens Refills _____
Directions: Inject 108mg subcutaneously every 2 weeks

☐ **OMVOH® (mirikizumab)** ☐ Prefilled Syringe ☐ Prefilled Pen
☐ UC INDUCTION DOSE: 300 mg IV at Week 0, Week 4, and Week 8 Qty: 1 vial Refills ___
☐ UC MAINTENANCE DOSE: 200mg SQ at Week 12 & every 4 weeks thereafter Qty: 1 pen/PFS Refills ___
☐ CD INDUCTION DOSE: 900 mg IV at Week 0, Week 4, and Week 8 Qty: 3 vials Refills ___
☐ CD MAINTENANCE DOSE: 300mg SQ at Week 12 & every 4 weeks thereafter Qty: 100mg + 200mg Refills ___

☐ **Rinvoq® (upadacitinib)**
☐ INDUCTION DOSE ☐ Take one 45mg tab once daily for 8 weeks Qty: 28 Refills one
☐ MAINTENANCE DOSE ☐ Take one 15mg tablet once daily Qty: 30 Refills _____
☐ Take one 30mg tablet once daily Qty: 30 Refills _____

☐ **Simponi® (golimumab)** TYPE: ☐ Smartject® Autoinjector ☐ Prefilled Syringe STRENGTH: ☐ 50mg ☐ 100mg
☐ 15 - 39kg Dosing & Directions: 100 mg SQ initially Week 0, 50 mg at Week 2, then 50 mg every 4 weeks Qty: QS 30 Days Refills _____
☐ ≥ 40kg Dosing & Directions: 200 mg SQ initially Week 0, 100 mg at Week 2, then 100 mg every 4 weeks Qty: QS 30 Days Refills _____

☐ **Skyrizi® (risankizumab-rzaa)** ☐ 600 mg single-dose vial - induction ☐ 180mg single dose cartridge ☐ 360mg single dose cartridge
☐ INITIATION: Infuse 600 mg as initial IV dose at Week 0, Week 4, and Week 8 ☐ Qty: 60 Days
☐ MAINTENANCE: 180 mg by SQ injection at week 12, and every 8 weeks thereafter ☐ Qty: 60 Days Refills _____
☐ MAINTENANCE: 360 mg by SQ injection at week 12, and every 8 weeks thereafter ☐ Qty: 60 Days Refills _____

☐ **Stelara® (ustekinumab)** ☐ 130mg Single Dose Vial ☐ 90mg single-dose prefilled syringe
☐ 260mg Starter (up to 55kg): Infuse intravenously over a period of at least one hour as directed
☐ 390mg Starter (greater than 55kg to 85kg): Infuse intravenously over a period of at least one hour as directed
☐ 520mg Starter (greater than 85mg): Infuse intravenously over a period of at least one hour as directed
☐ MAINTENANCE: Inject 90mg SQ every 8 weeks after initial intravenous dose 60 Days Supply? ☐YES ☐NO Refills _____

☐ **Tremfya® (guselkumab)** ☐ 100mg Pen ☐ 100mg One-Press Injector ☐ 100mg Prefilled Syringe ☐ 200mg Pen ☐ 200mg Prefilled Syringe
☐ INITIATION (Crohn's Only): Inject 400mg SQ at Week 0, Week 4 and Week 8 ☐ Qty: 60 Days
☐ MAINTENANCE: Inject 100mg SQ at Week 16, and every 8 weeks thereafter ☐ Qty: 60 Days Refills _____
☐ MAINTENANCE: Inject 200mg SQ at Week 12 and every 4 weeks thereafter ☐ Qty: 30 Days Refills _____

☐ **Xeljanz® (tofacitinib)**
☐ INDUCTION DOSE ☐ Take one 10mg tab twice daily for 8 weeks Qty: 120
☐ Take one 22mg XR tab once daily for 8 weeks Qty: 30
☐ MAINTENANCE DOSE ☐ Take one 5mg tablet twice daily Qty: 60 Refills _____
☐ Take one 10mg tablet twice daily Qty: 60 Refills _____
☐ Take one 11mg XR tablet once daily Qty: 30 Refills _____
☐ Take one 22mg XR tablet once daily Qty: 30 Refills _____

☐ **Zeposia® (ozanimod)** ☐ 7 Day Starter ☐ 0.92mg Capsule
☐ INDUCTION DOSE: Take 0.23 mg once daily for days 1-4, then take 0.46 mg once daily for days 5-7 Qty: 1 Pack
☐ MAINTENANCE DOSE: Take 0.92 mg once daily Qty: 30 Refills _____

CONFIDENTIALITY NOTICE: THE INFORMATION IN THIS TRANSMITTAL IS CONFIDENTIAL AND INTENDED ONLY FOR THE RECIPIENT LISTED ABOVE. IF YOU ARE NEITHER THE INTENDED RECIPIENT NOR A PERSON RESPONSIBLE FOR DELIVERING THIS TRANSMITTAL TO THE INTENDED RECIPIENT, YOU ARE HEREBY NOTIFIED THAT ANY DISTRIBUTION OR COPYING OF THIS TRANSMITTAL IS PROHIBITED. IF YOU RECEIVE THIS TRANSMITTAL IN ERROR, PLEASE IMMEDIATELY NOTIFY US AND RETURN THE TRANSMITTAL TO US AT OUR EXPENSE.